

TRANEXAMIC ACID (TXA) ADMINISTRATION**FOR USE IN ADULT PATIENTS (≥15 years old)****BLS**

Universal Protocol #601
 General Trauma Protocol #660
 Pulse Oximetry – O₂ administration per Airway Management Protocol #602

ALS Standing Orders

- **Indications:**
 - Blunt or penetrating traumatic injury with SBP ≤90mmHg
 - Significant blood loss with ongoing bleeding not controlled by direct pressure, hemostatic agents, or tourniquet application
- **Tranexamic Acid (TXA)** 1 gm in 100 mL IV infusion over 10 minutes, no repeat

Base Hospital Orders Only

As needed

Notes

Mixing the Concentration

- Maintain sterile technique
- 10mg/mL concentration
 - Supplies needed: 1 gm Tranexamic Acid (**TXA**), 100mL of **Normal Saline**
- Mixing instructions:
 - 1gm of **TXA** into 100mL **Normal Saline**
- Label the bag with the drug name and final concentration
 - Example: “TXA 1gm in 100mL”

Contraindications:

- >3 hours post injury
- Isolated extremity hemorrhage when bleeding has been controlled and if there is a strong expectation of re-implantation
- Isolated spinal shock
- Isolated head injury
- Non-blood loss conditions
- Traumatic Arrest with >5 min of CPR without ROSC
- Drowning or hanging victims
- <15 years old
- Active Thromboembolic event (within 24 hrs); i.e. CVA, MI, Pulmonary Embolism, or DVT
- Hypersensitivity or anaphylactic reaction to TXA

Possible adverse effects include:

- Hypotension with rapid IV infusion
- Chest tightness, difficulty breathing
- Facial flushing, swelling in hands or feet
- Blurred vision