County of San Luis Obispo Public Health Department Page 1 of 5

Division: Emergency Medical Services Agency Effective Date: 08/01/2018

POLICY #153: TRAUMA PATIENT TRIAGE AND DESTINATION

I. PURPOSE

A. To establish guidelines for EMS personnel to identify and transport "significantly injured" patients who could benefit from the rapid response and specialized services of a trauma center.

II. SCOPE

A. This policy applies to both adult and pediatric injured patients, unless stated otherwise.

III. PROCEDURE

- A. Trauma Activation Criteria
 - "STEP 1 or STEP 2 TRAUMA ALERT" Patient meeting <u>any one</u> of the Physiologic (Step 1) and/or Anatomic criteria (Step 2) following a traumatic event shall be designated a "TRAUMA ALERT" and transported to the closest trauma center.
 - "STEP 3 TRAUMA CONSULTATION" Patient meeting (Step 3) Mechanism of Injury - contact with the County of San Luis Obispo (SLO) Trauma Center for patient destination.
 - 3. "STEP 4 TRAUMA CONSULTATION"- shall be made with the SLO Trauma Center to determine destination when the paramedic identifies a significantly injured patient that <u>DOES NOT</u> meet the Step 1 (Physiologic), Step 2 (Anatomic) or Step 3 (Mechanism of Injury) criteria but meets one or more of the special patient or system considerations

B. Trauma Patient Criteria

Patients meeting any one of the Physiologic and/or Anatomic criteria following a traumatic event shall be a "TRAUMA ALERT" and transported to the closest trauma center. Patient meeting Mechanism of Injury and/or Special Patient/System Considerations shall be a TRUAMA CONSULT and contact the County of San Luis Obispo (SLO) Trauma Center for patient destination.

1. STEP 1 (Physiologic Criteria)

- a. **Adult** injured patients meeting any one of the following criteria:
 - 1. Glasgow Coma Scale ≤13 (based on patient history and attributed to injury)
 - 2. Systolic blood pressure <90 mmHg
 - 3. Respiratory rate <10 or >29 breaths per minute
- b. *Pediatric* injured patients (≤ 34 Kg) meeting any one of the following criteria:

- 1. Glasgow Coma Scale ≤13 (based on patient history and attributed to injury)
- 2. Evidence of poor perfusion color, temperature, etc.
- 3. Respiratory rate
 - >60 breaths per minute or respiratory distress
 - <20 breaths per minute in infants <1year

4. Heart rate

- ≤5 years (<22Kg) heart rate <80 beats per minute or >180 beats per minute
- ≥6 years (23-34Kg) heart rate <60 beats per minute or >160 beats per minute

5. Blood pressure

- Newborn (<1 month) systolic blood pressure <60mmHg
- Infant (1 month -1 year) systolic blood pressure <70mmHg
- Child (1 year-10 years) systolic blood pressure <70mmHg + 2X age in years
- Child (11-14 years) systolic blood pressure <90mmHg

2. STEP 2 (Anatomic Criteria)

Injured patients meeting any one of the following criteria:

- a. All significant penetrating injuries to head, neck, torso and extremities proximal to knee or elbow
- b. Chest wall instability or deformity (e.g. flail chest)
- c. Two proximal long bone fractures (above the elbows and or knees)
- d. Mangled, degloved or pulseless extremity
- e. Open or depressed skull fracture
- f. Paralysis
- g. Pelvic injury with high-risk mechanism of injury (motor vehicle collisions, auto vs. pedestrian accidents, motorcycle collisions, falls from heights)

3. STEP 3 (Mechanism of Injury Criteria)

Injured patients meeting <u>any one</u> of the following criteria:

- a. Falls
 - 1. Adults: >20 feet (one story is equal to 10 feet)
 - 2. Pediatric (≤ 34kg) : >10 feet or ≥ two times the height of the child
- b. High-risk auto crash:

- 1. Passenger Space Intrusion (PSI) of space: >12 inches occupant patient site; or >18 inches anywhere within the passenger space
- 2. Ejection (partial or complete) from automobile
- 3. Death in same passenger compartment
- c. Auto vs. pedestrian/bicyclist thrown, run over, or with significant impact (>20 mph)
- d. Motorcycle or unenclosed transport vehicle crash (>20 mph)

4. STEP 4 (Special Patient or System Considerations)

Age and co-morbid considerations.

- a. EMS provider judgment
- b. Age greater than 65
 - 1. SBP <110 mmHg may represent shock
- c. Pediatric (≤ 34kg)
- d. Pregnancy > 20 weeks
- e. Anticoagulation therapy (excluding aspirin) or other bleeding disorders with head injury (excluding minor injuries)
- f. Burns with trauma mechanism

Note:

A TRAUMA CONSULT <u>is not required</u> for ground level/low impact falls with GCS ≥ 14 or when the GCS is normal for patient

C. Contact the Trauma Center

Contact the receiving trauma center early and immediately upon determining the patient meets trauma patient triage criteria with a "TRAUMA ALERT" or "TRAUMA CONSULTATION"

1. "TRAUMA ALERT"

A "TRAUMA ALERT" is initiated when an injured patient meets any one of the Step 1 (Physiologic) or Step 2 (Anatomic) Criteria. Consider early notification to the intended receiving Trauma Center, from the scene when possible

- a. EMS personnel should provide a "TRAUMA ALERT" early and from the scene when possible to assist in early activation of the trauma team and determination of patient destination.
- b. ALS personnel must contact the trauma center with the TRAUMA ALERT.
- c. A "TRAUMA ALERT" report should include the following:
 - 1. "TRAUMA ALERT" meeting trauma triage step criteria "x"
 - 2. Unit and medic #

- ETA to trauma center
- 4. Report on individual patient (MIVT format):
 - Age and sex
 - Mechanism of injury
 - Injury and complaints
 - Vital signs including GCS
 - Treatment
 - Include <u>specific</u> triage findings or considerations that identify the patient as meeting TRAUMA ALERT criteria.

2. "TRAUMA CONSULTATION"

"TRAUMA CONSULTATION" with a SLO trauma center should be obtained to determine trauma patient destination when Step 3 (mechanism(s) of injury) criteria or Step 4 (special considerations) are present and Step 1 (physiologic) and Step 2 (anatomic) criteria are NOT met.

- a. Only ALS personnel may request a "TRAUMA CONSULTATION" for patient destination
- b. A "TRAUMA CONSULTATION" report should include the following:
 - 1. "TRAUMA CONSULTATION" meeting trauma triage step criteria "x"
 - 2. Unit and medic #
 - 3. ETA to trauma center and ETA to closest ED (When the trauma center is the closest facility include in the radio contact information notifying them they are the closest receiving facility)
 - 4. Report on the individual patient: (MIVT format)
 - Patient age and sex
 - Mechanism of injury and scene
 - Injury and complaints
 - Vital signs including GCS
 - Treatment and response
 - Include <u>specific</u> findings or considerations that identify the patient as meeting TRAUMA CONSULTATION criteria
- c. Paramedic Concerns
- 3. The Trauma center, when not receiving the patient, shall notify the receiving hospital of the incoming patient and provide that hospital with the prehospital care patient information.
- 4. When practical, a brief updated report should be given to the trauma center Hospital and include any significant changes in route in vital signs, GCS, physical findings, symptoms or treatments.

D. Exceptions to Direct Transport to a Trauma Center

Trauma patients will be transported to the closest ED in the following situations:

- 1. Patient condition necessitates transport to the closest ED, such as the following:
 - a. Unmanageable airway (intubation attempts are unsuccessful and an adequate airway cannot be maintained with BVM or other device)
 - b. Uncontrollable bleeding with rapidly deteriorating vital signs
 - c. Traumatic cardiac arrest see EMS Agency Prehospital Determination of Death/Do Not Resuscitate (DNR) End of Life Care Policy #125.
- 2. SLO Trauma Center destination order
- 3. Patient refusal see EMS Agency Patient Refusal of Treatment and/or Transport Policy #203.
- 4. Trauma center is on complete diversion see EMS Agency Hospital Diversion Policy #154: Hospital Diversion.
- E. The utilization of EMS helicopter for the response and transport of trauma patients must be in accordance with EMS Agency EMS Helicopter Operations. EMS Helicopter Policy #155 transport should be considered when ground transport is greater than 30 minutes from the trauma center and air transport would be more expeditious than ground transport.

IV. AUTHORITY

- California Health and Safety Code, Division 2.5.
- California Code of Regulations, Title 22, Chapter 7

V. ATTACHMENTS

A. Trauma Triage Matrix