

UNIVERSAL		
MEDICAL	TRAUMA	
BLS		
<ul style="list-style-type: none"> Evaluate Scene Safety/Personal Protective Equipment Assess, establish and maintain airway <ul style="list-style-type: none"> Suction as needed Pulse Oximetry <ul style="list-style-type: none"> O₂ administration per Airway Management Protocol #602 Evaluate breathing and circulation Assess chief complaint Focused physical exam and vital signs: <ul style="list-style-type: none"> Pulse Blood pressure Respiratory rate Lung sounds Skin signs BLS treatment protocols Optional skills as approved by SLOEMSA. 	<ul style="list-style-type: none"> Evaluate Scene Safety/Personal Protective Equipment Assess, establish and maintain airway <ul style="list-style-type: none"> Suction as needed Pulse Oximetry <ul style="list-style-type: none"> O₂ administration per Airway Management Protocol #602 Evaluate breathing and circulation Control life-threatening bleeding Remove patient's clothing to expose and identify injuries Ensure patient warmth – cover patient after clothing removal to maintain core body temperature Spinal motion restriction (SMR) if indicated per Spinal Motion Restriction Procedure # 702 BLS treatment protocols Optional skills as approved by SLOEMSA 	
ALS		
<ul style="list-style-type: none"> Vascular access – Procedure #710 Consider 12-lead ECG early Capnography (if available/applicable) Blood Glucose Measurement Transport Determination ALS Treatment Protocols <ul style="list-style-type: none"> Adult Consider Normal Saline up to 500mL IV <ul style="list-style-type: none"> May repeat x1 for persistent hypotension Pediatric Consider Normal Saline up to 10mL/kg <ul style="list-style-type: none"> May repeat x1 for persistent hypotension 	<ul style="list-style-type: none"> Trauma Triage and Destination ALS Treatment Protocols 	
Base Hospital Orders Only		
<ul style="list-style-type: none"> Determined on patient needs If applicable, see Policy #219: Assisting Patients with Their Emergency Medications 	<ul style="list-style-type: none"> Determined on patient needs 	
Notes		
<ul style="list-style-type: none"> Use Pediatric Policies for patients ≤34 kg and consider use of Broselow tape or equivalent Rapid transport for Specialty Care patients (Trauma, STEMI, CVA-TIA). Target scene departure ≤ 10 minutes for transport personnel. 		