

ALLERGIC REACTION/ANAPHYLAXIS	
ADULT	PEDIATRIC (≤34 KG)
BLS	
<ul style="list-style-type: none"> • Universal Protocol #601 • Pulse Oximetry <ul style="list-style-type: none"> ○ O₂ administration per Airway Management Protocol #602 • May assist with the administration of patient’s prescribed medication (i.e. Epi Auto-injector, inhaler, etc.) 	<p>Same as Adult</p>
BLS Elective Skill (Approved Providers Only)	
<p>Unstable (Dyspnea/Wheezing/Shock)</p>	<p>Unstable (Dyspnea/Wheezing/Shock)</p>
<ul style="list-style-type: none"> • Adult 0.3 mg Epinephrine Auto-Injector administered in anterolateral thigh <ul style="list-style-type: none"> ○ May repeat, if indicated, every 5 min, max 3 doses 	<ul style="list-style-type: none"> • Pediatric (≥15 kg) 0.15 mg Epinephrine Auto-Injector administered in anterolateral thigh <ul style="list-style-type: none"> ○ May repeat, if indicated, every 5 min, max 3 doses
BLS Optional Scope Skill (Approved Providers Only)	
<p>Unstable (Dyspnea/Wheezing/Shock)</p>	<p>Unstable (Dyspnea/Wheezing/Shock)</p>
<ul style="list-style-type: none"> • Adult Epinephrine 1:1000 0.3 mg IM <ul style="list-style-type: none"> ○ May repeat, if indicated, every 5 min, max 3 doses 	<ul style="list-style-type: none"> • Pediatric (≥15 kg), Epinephrine 1:1000 0.15 mg IM anterolateral thigh <ul style="list-style-type: none"> ○ May repeat, if indicated, every 5 min , max 3 doses
ALS Standing Orders	
<p>Stable (Itching/rash)</p>	<p>Stable (Itching/rash)</p>
<ul style="list-style-type: none"> • Diphenhydramine 50 mg IV/IM 	<ul style="list-style-type: none"> • Diphenhydramine 2 mg/kg IV/IM – not to exceed 50 mg
<p>Unstable (Dyspnea/Wheezing/Shock)</p>	<p>Unstable (Dyspnea/Wheezing/Shock)</p>
<ul style="list-style-type: none"> • Albuterol 2.5-5 mg via HHN/Mask/BVM with adjunct, over 5-10 min <ul style="list-style-type: none"> ○ repeat as needed • Epinephrine 1:1,000 0.01 mg/kg IM – not to exceed 0.5 mg <ul style="list-style-type: none"> ○ may repeat every 5 min, max 3 doses • Diphenhydramine 50 mg IV/IM 	<ul style="list-style-type: none"> • Albuterol 2.5-5 mg via HHN/Mask/BVM with adjunct, over 5-10 min <ul style="list-style-type: none"> ○ repeat as needed • Epinephrine 1:1,000 0.01 mg/kg IM – not to exceed 0.3 mg <ul style="list-style-type: none"> ○ may repeat every 5 min, max 3 doses • Diphenhydramine 2 mg/kg IV/IM – not to exceed 50 mg
<p>Extremis</p>	<p>Extremis</p>
<ul style="list-style-type: none"> • Epinephrine 1:1,000 0.01 mg/kg SL – not to exceed 0.5 mg <ul style="list-style-type: none"> ○ may repeat every 5 min, max 3 doses 	<ul style="list-style-type: none"> • Epinephrine 1:1,000 0.01 mg/kg SL – not to exceed 0.3 mg <ul style="list-style-type: none"> ○ may repeat every 5 min, max 3 doses

Base Hospital Orders Only	
Unresponsive to previous therapy <ul style="list-style-type: none"> • Epinephrine 1:10,000 0.01 mg/kg slow IV titrated – not to exceed 0.5 mg • As needed 	Unresponsive to previous therapy <ul style="list-style-type: none"> • Epinephrine 1:10,000 0.01 mg/kg slow IV titrated – not to exceed 0.3 mg • As needed
Notes	
<ul style="list-style-type: none"> • Auto-injector injection site should be exposed and cleansed with aseptic technique prior to injection. • Follow manufacturer’s instructions when using Epinephrine auto-injector. 	